



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## GENERAL NOTES.

**The Roseate Tern (*Sterna dougalli*) on Lake Michigan.**—On August 14, 1916, while watching the large number of terns congregated on the extreme southern end of Lake Michigan near Millers, Ind., an individual was noticed standing alone at the water's edge, which on being examined with field glasses, looked different from either the Forster's or the Common Tern, both of which were there in abundance. The specimen was collected, and proved to be an adult male Roseate Tern, in full breeding plumage. While the beautiful rosy tint on the breast was evident enough with the specimen in hand, it was not noticed while watching the bird on the beach. While this appears to be the first Lake Michigan record, it is not unlikely that careful watching would show an occasional wanderer of this species among the large number of terns that frequent this locality in fall. The skin is in the Harris Extension collection.—H. L. STODDARD, N. W. Harris Public School Extension of Field Museum, Chicago, Ill.

**Relationship of Florida Herons.**—I note in the October number of 'The Auk' which has just come to hand, on page 431 the statement that *Ardea herodias wardi* and *Ardea herodias wurdemanni* are both to be considered phases of *Ardea herodias occidentalis*.

My experience with the birds on the Florida Keys for the last five years leads me to believe that this dictum should not be adopted.

The Great White Heron of the Keys is so entirely different in its habits and psychological manifestation from the Great Blue Heron which occupies the same region that no one who knows the two birds in the field would believe that they were the same. The Great White Heron is of more social habits than the Blue Heron. You frequently see small groups of this species in a confined space. For example: on Duck Key, a small island less than sixty yards across, I found four of the Great White Herons. Then again there is a breeding colony on a small island in the lagoon on Chase's Key, which is used as a breeding ground exclusively by the Great White Heron.

The Great Blue Heron is more sparingly represented, and lacks the social habits of the white bird; that is, when not on its breeding ground.

I think that Mr. Oberholser's dictum is the correct one, and we should reserve the name *Ardea occidentalis* for the Great White Heron of the Florida Keys, and *Ardea herodias wardi* for the Great Blue Heron of that region.—PAUL BARTSCH, U. S. National Museum, Washington, D. C.

**A New Record for New England.**—This museum has obtained a specimen of the Mountain Plover (*Podasocys montanus* (Townsend)) taken at Chatham, Mass. It is an immature male and was shot on October 28 by Mr. A. E. Crowell. This bird was associated with some Black-breasted Plovers at the time.—W. SPRAGUE BROOKS, Boston Society of Natural History.